



Korean treatment recommendations for patients with axial spondyloarthritis

Mi Ryoung Seo¹, Jina Yeo¹, Jun Won Park², Yeon-Ah Lee³, Ju Ho Lee⁴, Eun Ha Kang⁴, Seon Mi Ji⁵, Seong-Ryul Kwon⁶, Seong-Kyu Kim⁷, Tae-Jong Kim⁸, Tae-Hwan Kim⁹, Hye Won Kim¹⁰, Min-Chan Park¹¹, Kichul Shin¹², Sang-Hoon Lee¹³, Eun Young Lee², Hoon Suk Cha¹⁴, Seung Cheol Shim¹⁵, Youngim Yoon², Seung Ho Lee¹⁶, Jun Hong Lim¹⁷, Han Joo Baek¹, and on behalf of the Korean Society of Spondyloarthritis Research

¹Division of Rheumatology, Department of Internal Medicine, Gil Medical Center, Gachon University College of Medicine, Incheon; ²Division of Rheumatology, Department of Internal Medicine, Seoul National University College of Medicine, Seoul; ³Division of Rheumatology, Department of Internal Medicine, Kyung Hee University College of Medicine, Seoul; ⁴Division of Rheumatology, Department of Internal Medicine, Seoul National University Bundang Hospital, Seongnam; ⁵National Health Insurance Service, Wonju; ⁶Division of Rheumatology, Department of Internal Medicine, Inha University College of Medicine, Incheon; ⁷Division of Rheumatology, Department of Internal Medicine, Daegu Catholic University School of Medicine, Daegu; ⁸Division of Rheumatology, Department of Internal Medicine, Chonnam National University Medical School and Hospital, Gwangju; ⁹Department of Rheumatology, Hanyang University Hospital for Rheumatic Diseases, Seoul; ¹⁰Division of General Internal Medicine, Department of Internal Medicine, Seoul National University Bundang Hospital, Seongnam; ¹¹Division of Rheumatology, Department of Internal Medicine, Yonsei University College of Medicine, Seoul; ¹²Division of Rheumatology, Department of Internal Medicine, Seoul Metropolitan Government-Seoul National University Boramae Medical Center, Seoul; ¹³Department of Rheumatology, Kyung Hee University Hospital at Gangdong, Kyung Hee University College of Medicine, Seoul; ¹⁴Division of Rheumatology, Department of Internal Medicine, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul; ¹⁵Division of Rheumatology, Daejeon Rheumatoid & Degenerative Arthritis Center, Chungnam National University Hospital, Daejeon; ¹⁶Korea Ankylosing Spondylitis Society, Seoul; ¹⁷Korea Ankylosing Spondylitis Corporation, Daejeon, Korea

Korean J Intern Med 2023;38:620-640
<https://doi.org/10.3904/kjim.2023.194>

In the article “Korean treatment recommendations for patients with axial spondyloarthritis” by Seo MR et al. (Korean J Intern Med 2023;38:620-640), there is a mistake in the first author’s affiliation. The affiliation 18 should be added and corrected as follows:

BEFORE

Mi Ryoung Seo¹

¹Division of Rheumatology, Department of Internal Medicine, Gil Medical Center, Gachon University College of Medicine, Incheon, Korea

AFTER

Mi Ryoung Seo^{1,18}

¹Division of Rheumatology, Department of Internal Medicine, Gil Medical Center, Gachon University College of Medicine, Incheon, Korea

¹⁸Department of Medicine, Inha University Graduate School, Incheon, Korea

We apologize for any inconvenience that this may have caused.