

Supplementary Table 2. The prevalence of high-risk OSA for measures of hypertension, diabetes, and obesity, according to decile group

Decile	Prevalence of high-risk OSA according to decile group											
	SBP		DBP		FPG		HbA1c		BMI		WC	
	%	SE	%	SE	%	SE	%	SE	%	SE	%	SE
1	4.2	1.0	7.7	1.1	6.8	1.2	7.1	1.2	1.3	0.4	1.7	0.6
2	7.2	1.0	6.8	1.0	5.1	0.8	5.4	0.8	2.9	0.8	5.1	1.0
3	7.5	1.2	6.8	1.0	7.0	1.0	9.3	1.3	4.8	0.9	9.8	1.2
4	9.8	1.3	9.2	1.3	11.8	1.8	13.9	1.6	7.5	1.2	9.9	1.3
5	13.8	1.5	11.3	1.6	12.2	1.4	10.3	1.3	12.4	1.5	9.9	1.2
6	17.2	1.8	12.1	1.3	13.5	1.6	12.7	1.5	10.2	1.2	12.8	1.4
7	14.7	1.5	11.5	1.4	13.8	1.5	14.0	1.8	18.3	1.6	12.2	1.3
8	18.9	1.8	14.7	1.4	16.1	1.7	16.3	1.3	12.6	1.5	13.9	1.6
9	22.0	1.7	15.8	1.6	20.8	1.8	21.7	2.1	22.4	2.0	20.4	1.9
10	19.0	1.6	30.9	1.9	25.7	1.9	24.4	1.8	35.8	2.2	37.2	2.1

OSA, obstructive sleep apnea; SBP, systolic blood pressure; DBP, diastolic blood pressure; FPG, fasting plasma glucose; HbA1c, glycosylated hemoglobin; BMI, body mass index; WC, waist circumference; SE, standard error.