

Supplementary Table1. Characteristics of antibiotics administered within 30 days

Variable	MDR bacteremia (n = 17)	Non MDR bacteremia (n = 33)	p value
Classification			
1st or 2nd generation cephalosporin	0 (0.0)	3 (9.1)	0.54
3rd or 4th generation cephalosporin	5 (29.4)	9 (27.3)	> 0.999
Fluoroquinolones	2 (11.8)	3 (9.1)	> 0.999
Amoxicillin or amoxicillin/clavulanate	2 (11.8)	1 (3.0)	0.24
Piperacillin/tazobactam	3 (17.6)	11 (33.3)	0.47
Carbapenem	3 (17.6)	1 (3.0)	0.09
Others ^{a)}	5 (29.5)	16 (48.5)	0.39
Unknown	2 (11.8)	2 (6.1)	0.60
Route			
Intravenous	4 (23.5)	9 (27.3)	> 0.999
Oral	8 (47.1)	13 (39.4)	0.60
Both (intravenous + oral)	3 (14.6)	9 (27.3)	0.51
Unknown	2 (11.8)	2 (6.1)	0.60
Duration			
≤ 1 week	10 (58.8)	23 (69.7)	0.44
> 1 week	7 (41.2)	10 (30.3)	0.44

Values are presented as number (%).

MDR, multidrug-resistant.

^{a)}Others included trimethoprim-sulfamethoxazole, rifamycin, azithromycin, and metronidazole.