

**Supplementary Table 1. Predictive factors for 30-day mortality in patients with low risk endoscopic stigmata (n = 66)**

Factor	Univariate analysis			Multivariate analysis	
	30-Day death (+) (n = 5)	30-Day death (-) (n = 61)	p value	Odds ratio (95% CI)	p value
Male sex	3 (60.0)	45 (73.8)	0.608	-	-
Age ≥ 65 yr	5 (100.0)	28 (45.9)	0.053	-	-
Initial vital signs					
SBP < 90 mmHg	0	2 (3.3)	1.000	-	-
Heart rate > 100 beat/min	0	15 (24.6)	0.581	-	-
Initial laboratory data					
Hemoglobin, g/dL	6.40 ± 2.89	8.04 ± 2.56	0.177	-	-
Platelets, × 10 <sup>9</sup> /L	248 ± 178	246 ± 113	0.965	-	-
Blood nitrogen urea, mg/dL	84.60 ± 31.92	40.64 ± 29.19	0.002 <sup>a</sup>	1.03 (0.99–1.08)	0.174
Prothrombin time, sec	16.02 ± 10.32	12.43 ± 3.86	0.481	-	-
Severe bleeding related symptoms (hematemesis, hematochezia)	2 (40.0)	25 (41.0)	1.000	-	-
Positive nasogastric tube aspiration <sup>b</sup>	3 (60.0)	29 (47.5)	1.000	-	-
Positive rectal examination <sup>c</sup>	5 (100.0)	47 (77.0)	0.576	-	-
Heavy alcoholics	2 (40.0)	23 (37.7)	1.000	-	-
Current smoker	1 (20.0)	23 (37.7)	0.645	-	-
Past history of gastrointestinal bleeding	0	17 (27.9)	0.317	-	-
Past history of peptic ulcer disease	1 (20.0)	19 (31.1)	1.000	-	-
Transfusion requirement	5 (100.0)	40 (65.6)	0.169	-	-
Red cell transfusion unit	2.60 ± 0.89	1.72 ± 1.69	0.258	-	-
Comorbidities					
Hypertension	4 (80.0)	30 (49.2)	0.357	-	-
Diabetes mellitus	5 (100.0)	17 (27.9)	0.003 <sup>a</sup>	24.79 (0.12–81.98)	0.107
Cardiovascular disease	1 (20.0)	13 (21.3)	1.000	-	-
Liver cirrhosis	0	14 (23.0)	0.576	-	-
Chronic kidney disease	2 (40.0)	6 (9.8)	0.047 <sup>a</sup>	2.34 (0.82–66.62)	0.619
Cerebrovascular disease	2 (40.0)	8 (13.1)	0.162	-	-
Heart failure	3 (60.0)	11 (18.0)	0.027 <sup>a</sup>	2.81 (0.29–27.21)	0.373
Metastatic malignancy	0	3 (4.9)	1.000	-	-
Peripheral vascular disease	1 (20.0)	1 (1.6)	0.147	-	-
Medication					
Antiplatelet agents	2 (40.0)	22 (36.1)	1.000	-	-
Vitamin K antagonist	1 (20.0)	0	0.076	-	-
Nonsteroidal anti-inflammatory drug	2 (40.0)	13 (21.3)	0.318	-	-
PPI comedication	0	2 (3.3)	1.000	-	-
Blatchford score ≥ 12	5 (100.0)	43 (70.5)	0.312	-	-
Rockall score ≥ 5	4 (80.0)	25 (41.0)	0.160	-	-
Endoscopy lesion location	-	-	1.000	-	-
Body	1 (20.0)	10 (16.4)	-	-	-
Other than body	4 (80.0)	51 (83.6)	-	-	-
Tachycardia (heart rate > 100 beat/min) during the admission	2 (40.0)	17 (27.9)	0.621	-	-
Hypotension (SBP < 90 mmHg) during the admission	2 (40.0)	6 (9.8)	0.107	-	-
Rebleeding within 30 days	1 (20.0)	2 (3.3)	0.214	-	-
Hospital stay, day	9.80 ± 8.44	4.82 ± 4.95	0.259	-	-

Values are presented as number (%) or mean ± SD.

CI, confidence interval; SBP, systolic blood pressure; PPI, proton pump inhibitor.

<sup>a</sup>Significant values with p < 0.05.<sup>b</sup>Nasogastric tube was performed in 64 patients.<sup>c</sup>Rectal examination was performed in 66 patients.