

Supplementary Table 1. Multivariable Cox proportional hazards model for heart disease and stroke by levels of Lp(a)

Variable	Hazard ratio (95% CI)			
	Model 1	Model 2	Model 3	Model 4
Stroke (n = 97/833, 11.6%)				
Lp(a)				
1st Quartile	1.00	1.00	1.00	1.00
2nd Quartile	0.80 (0.41–1.58)	0.78 (0.39–1.54)	0.76 (0.38–1.51)	0.81 (0.41–1.62)
3rd Quartile	1.08 (0.57–2.04)	0.98 (0.51–1.88)	0.98 (0.51–1.88)	1.03 (0.54–1.98)
4th Quartile	2.11 (1.19–3.75)	1.01 (1.22–3.27)	1.86 (1.03–3.36)	2.10 (1.16–3.80)
p for trend				< 0.001
Heart disease (n = 114/833, 13.7%)				
Lp(a)				
1st Quartile	1.00	1.00	1.00	1.00
2nd Quartile	0.77 (0.40–1.48)	0.74 (0.38–1.43)	0.73 (0.38–1.42)	0.78 (0.40–1.53)
3rd Quartile	1.96 (1.13–3.41)	1.85 (1.06–3.25)	1.84 (1.04–3.23)	1.96 (1.11–3.46)
4th Quartile	2.01 (1.15–3.52)	1.70 (0.96–3.00)	1.72 (0.97–3.06)	1.85 (1.04–3.30)
p for trend				< 0.001

Lp(a), lipoprotein(a); CI, confidence interval.