

Supplementary Table 3. Comparison of the FPG and 2hr PG criteria for the diagnosis of diabetes according to BMI categories

FPG, mg/dL	2hr PG, mg/dL				DM, %
	< 140	140–199	≥ 200	Total	
23.0 ≤ BMI < 25.0 kg/m ² (n = 604)					
< 100	28.6	15.9	2.8	47.4	5.9
100–109	12.7	11.9	5.3	30.0	17.7
110–125	3.5	7.8	6.3	17.5	35.8
≥ 126	0.2	0.8	4.1	5.1	100
25.0 ≤ BMI < 27.0 kg/m ² (n = 487)					
< 100	22.6	9.9	6.0	38.4	15.5
100–109	11.3	12.9	7.4	31.6	23.4
110–125	4.1	8.8	10.5	23.4	44.7
≥ 126	0.5	1.2	5.1	6.6	100
BMI ≥ 27.0 kg/m ² (n = 546)					
< 100	19.4	12.5	3.1	35.0	8.9
100–109	13.4	14.1	5.5	33.0	16.7
110–125	3.1	9.3	11.7	24.1	74.4
≥ 126	0.9	1.1	5.9	7.9	100

FPG, fasting plasma glucose; 2hr PG, 2-hour postchallenge glucose; BMI, body mass index, DM, diabetes mellitus.