

Supplementary Table 2. Age and sex profiles of the incident cases of NTM-PD and incidence rate (per 100,000 person-year) during the study period

| Age group, yr | Incident cases of NTM-PD | | | | Rate ratio ^a | Total number | |
|------------------|--------------------------|-------------------------|------------|-------------------------|----------------------------|--------------|------------|
| | Male | | Female | | | Male | Female |
| | No. (%) | Incidence rate (95% CI) | No. (%) | Incidence rate (95% CI) | | | |
| 30–39 | 5 (1.2) | 0.5 (0.2–1.1) | 19 (3.2) | 1.1 (0.7–1.8) | 2.4 | 1,081,209 | 1,704,841 |
| 40–49 | 13 (3.1) | 0.9 (0.5–1.6) | 74 (12.4) | 3.5 (2.8–4.5) | 3.9 | 1,445,775 | 2,091,878 |
| 50–59 | 69 (16.5) | 3.2 (2.5–4.1) | 207 (34.6) | 7.2 (6.3–8.3) | 2.2 | 2,129,300 | 2,869,038 |
| 60–69 | 143 (34.1) | 6.1 (5.2–7.2) | 158 (26.4) | 6.2 (5.3–7.3) | 1.0 | 2,338,416 | 2,531,907 |
| 70–79 | 150 (35.8) | 9.1 (7.7–10.7) | 108 (18.1) | 6.8 (5.6–8.2) | 0.7 | 1,654,679 | 1,596,645 |
| ≥ 80 | 36 (8.6) | 9.5 (6.7–13.2) | 28 (4.7) | 7.5 (5.0–10.9) | 0.8 | 380,805 | 375,772 |
| Total | 419 (100) | 3.3 (3.0–3.6) | 598 (100) | 4.0 (3.7–4.4) | 1.2 | 12,798,235 | 14,829,610 |

NTM-PD, nontuberculous mycobacterial pulmonary disease; CI, confidence interval.

^aRatio of female to male incidence rate.