

Supplementary Table 2. Pathologic diagnosis in 1,932 NDRD patients

Pathologic diagnosis	Number	% among NDRD
IGAN	473	23.1
MN	296	14.5
FSGS	226	11.0
MCD	153	7.5
MPGN	86	4.2
NONSPGN	85	4.2
CRESGN	82	4.0
ATIN	81	4.0
SCLEROSIS	80	3.9
ATN	76	3.7
HT-N	73	3.6
CTIN	47	2.3
LN	42	2.1
INADEQUATE	28	1.4
AMYLODIOSIS	22	1.1
PIGN	22	1.1
HSN	19	0.9
NORMAL	18	0.9
THIN	18	0.9
TIN	18	0.9
TMA	18	0.9
VASCULITIS	18	0.9
ACTIN	17	0.8
MYELOMA	16	0.8
LCDD	5	0.2
C ₃ G	4	0.2
IGMN	4	0.2
ANTIGBM	3	0.1
C ₁ Q	3	0.1
LC-N	3	0.1
APN	2	0.1
FIBRILLARY	2	0.1
ALPORT	1	0.0
ANALGESIC	1	0.0
CPN	1	0.0
CRYO	1	0.0
ECD	1	0.0
ISCHEMIC	1	0.0
HFRS	1	0.0
Number of diagnosis	2,047	
Number of patients	1,932	
Number of patients with two diagnosis	115	

NDRD, non-diabetic renal disease; IGAN, IgA nephropathy; MN, membranous nephropathy; FSGS, focal segmental glomerulosclerosis; MCD, minimal change disease; MPGN, membranoproliferative glomerulonephritis; NONSPGN, non-specific glomerulonephritis; CRESGN, crescentic glomerulonephritis; ATIN, acute tubulointerstitial nephritis; ATN, acute tubular necrosis; HT-N, hypertensive nephrosclerosis; CTIN, chronic tubulointerstitial nephritis; LN, lupus nephritis; PIGN, post-infectious glomerulonephritis; HSN, Henoch-Schonlein nephritis; THIN, thin membrane disease; TIN, tubulointerstitial nephritis; TMA, thrombotic microangiopathy; ACTIN, acute and chronic tubulointerstitial nephritis; LCDD, light chain deposition disease; C₃G, C₃ glomerulopathy; IGMN, IgM nephropathy; GBM, glomerular basement membrane; LC-N, liver cirrhosis-related nephropathy; APN, acute pyelonephritis; CPN, chronic pyelonephritis; CRYO, cryoglobulinemic glomerulonephritis; ECD, Erdheim-chester disease; HFRS, hemorrhagic fever with renal syndrome.