

Supplementary Table 4. Risk factors of ESKD in adult diabetic patients.

Variable	B	HR	95% CI	p value
Male sex	0.21	1.23	1.02–1.48	0.03
Age, yr	-0.02	0.98	0.98–0.99	< 0.001
Hypertension (presence)	0.37	1.45	1.08–1.96	0.01
SBP (compared to 120–129 mmHg)				
< 110	0.13	1.14	0.75–1.72	0.55
110–119	0.09	1.10	0.78–1.55	0.61
130–139	0.52	1.68	1.23–2.30	0.001
140–149	0.30	1.35	0.99–1.87	0.06
≥ 150	0.31	1.36	1.00–1.85	< 0.05
eGFR, mL/min/1.73 m ²	-0.03	0.97	0.97–0.98	< 0.001
Serum albumin, g/dL	-0.21	0.81	0.70–0.93	0.004
Hemoglobin, g/dL	-0.13	0.88	0.84–0.92	< 0.001
Pathology (compared to NDRD)				
NDRD/DN	0.01	1.01	0.58–1.78	0.96
Pure DN	0.57	1.76	1.46–2.12	< 0.001

Model adjusted with age, gender, presence of hypertension, SBP, diastolic blood pressure, serum albumin, hemoglobin, glomerular filtration rate, and urine protein-to-creatinine ratio.

ESKD, end-stage kidney disease; B, beta coefficient; HR, hazard ratio; CI, confidence interval; SBP, systolic blood pressure; eGFR, estimated glomerular filtration rate; NDRD, non-diabetic renal disease; DN, diabetic nephropathy.