

Supplementary Table 5. Diagnostic indices of the clinical criteria of HT-N to distinguish biopsy-proven HT-N in total cohort.

Clinical criteria	No. of biopsy-proven HT-N	Sensitivity (95% CI)	Specificity (95% CI)	PPV (95% CI)	NPV (95% CI)	DOR (95% CI)
Criteria 1 (n = 10,659)	368	1.000 (0.990–1.000)	0.466 (0.459–0.473)	0.035 (0.031–0.038)	1.000 (1.000–1.000)	-
Criteria 2 (n = 6,839)	253	0.688 (0.637–0.735)	0.668 (0.662–0.675)	0.037 (0.033–0.042)	0.991 (0.990–0.993)	4.429 (3.546–5.532)
Criteria 3 (n = 2,927)	142	0.386 (0.336–0.438)	0.860 (0.855–0.864)	0.049 (0.041–0.057)	0.987 (0.985–0.989)	3.849 (3.108–4.766)

Criteria 1 includes only hypertension. Criteria 2 includes hypertension, no diabetes, and no evidence of vasculitis or lupus nephritis. Criteria 3 is criteria 2 plus urine protein-to-creatinine ratio < 2.0 g/g.

HT-N, hypertensive nephrosclerosis; CI, confidence interval; PPV, positive predictive value; NPV, negative predictive value; DOR, diagnostic odds ratio.