

Supplementary Table 6. Pathologic diagnosis in 2,927 patients with clinical HT-N

	Biopsy-proven HT-N	Non-HT-N ^a	Sum
IGAN	17 (9.4)	1,566 (54.4)	1,583 (51.7)
MN	3 (1.7)	143 (5.0)	146 (4.8)
FSGS	4 (2.2)	288 (10.0)	292 (9.5)
MCD	0	143 (5.0)	143 (4.7)
DN	0	0	0
LN	0	0	0
MPGN	1 (5.5)	59 (2.0)	60 (2.0)
HT-N	142 (78.5)	0	142 (4.6)
NONSPFGN	2 (1.1)	125 (4.3)	127 (4.1)
SCLEROSIS	0	65 (2.3)	65 (2.1)
CRESGN	0	25 (0.9)	25 (0.8)
ATN	5 (2.8)	57 (2.0)	62 (2.0)
ATIN	1 (5.5)	54 (1.9)	55 (1.8)
CTIN	0 (0)	61 (2.1)	61 (2.0)
NORMAL	2 (1.1)	52 (2.1)	54 (2.0)
THIN	0	63 (2.2)	63 (2.1)
INADEQUATE	1 (5.5)	24 (0.8)	25 (0.8)
PIGN	0	18 (0.6)	18 (0.6)
HSN	0	22 (0.8)	22 (0.7)
AMYLODIOSIS	1 (5.5)	6 (0.2)	7 (0.2)
TIN	1 (5.5)	22 (0.8)	23 (0.8)
TMA	1 (5.5)	22 (0.8)	23 (0.8)
VASCULITIS	0	5 (0.2)	5 (0.2)
HFRS	0	6 (0.2)	6 (0.2)
MYELOMA	0	3 (0.1)	3 (0.1)
ACTIN	0	8 (0.3)	8 (0.3)
C3G	0	7 (0.2)	7 (0.2)
C1Q	0	9 (0.3)	9 (0.3)
LIGHT CAHIN	0	4 (0.1)	4 (0.1)
IGMN	0	4 (0.1)	4 (0.1)
ALPORT	0	4 (0.1)	4 (0.1)
ANTIGBM	0	0	0
LC-N	0	0	0
HUS	0	3 (0.1)	3 (0.1)
GBM	0	4 (0.1)	4 (0.1)
CPN	0	2 (0.1)	2 (0.1)
CORTICALNECROSIS	0	2 (0.1)	2 (0.1)
ISCHEMIC NEPHROPATHY	0	0	0
ANALGESIC	0	2 (0.1)	2 (0.1)
APN	0	1 (0.03)	1 (0.03)
FABRY	0	0	0
FIBRILLARY	0	0	0

Supplementary Table 6. Continued

	Biopsy-proven HT-N	Non-HT-N ^a	Sum
CRYO	0	0	0
CRYSTAL	0	0	0
GRANULOMA	0	0	0
ECD	0	0	0
NECGN	0	0	0
TTP	0	1 (0.03)	1 (0.03)
WM	0	1 (0.03)	1 (0.03)
Sum of diagnosis	181	2,881	3,062
No. of patients	142	2,785	2,927
No. of patients with two diagnosis	39	96	135

Values are presented as number (%).

HT-N, hypertensive nephrosclerosis; IGAN, IgA nephropathy; MN, membranous nephropathy; FSGS, focal segmental glomerulosclerosis; MCD, minimal change disease; DN, diabetic nephropathy; LN, lupus nephritis; MPGN, membranoproliferative glomerulonephritis; NONSPFGN, non-specific glomerulonephritis; CRESGN, crescentic glomerulonephritis; ATN, acute tubular necrosis; ATIN, acute tubulointerstitial nephritis; CTIN, chronic tubulointerstitial nephritis; THIN, thin membrane disease; PIGN, post-infectious glomerulonephritis; HSN, Henoch-Schonlein nephritis; TIN, tubulointerstitial nephritis; TMA, thrombotic microangiopathy; HFRS, hemorrhagic fever with renal syndrome; ACTIN, acute and chronic tubulointerstitial nephritis; C3G, C3 glomerulopathy; IGMN, IgM nephropathy; GBM, glomerular basement membrane; LC-N, ; HUS, hemolytic uremic syndrome; GBM, non-specified glomerular basement membrane abnormality; CPN, chronic pyelonephritis; APN, acute pyelonephritis; CRYO, cryoglobulinemic glomerulonephritis; ECD, Erdheim-Chester disease; NECGN, necrotizing glomerulonephritis; TTP, thrombotic thrombocytopenic purpura; WM, Waldenström's macroglobulinemia.

^aPathologic diagnoses other than HT-N.