

Supplementary Table 9. Clinical characteristics of matched cohort among patients with biopsy-proven HT-N or non-HT-N

Characteristic	Non-HT-N	Biopsy-proven HT-N	p value
Number	206	103	
Age, yr	44.5 ± 13.5	44.7 ± 13.7	0.897
Male sex	149 (72.3)	70 (68.0)	0.507
Period of biopsy			
1979–1989	-	-	
1990–1999	15 (7.3)	8 (7.8)	
2000–2009	46 (22.3)	22 (21.4)	
2010–2018	145 (70.4)	73 (70.9)	
SBP, mmHg	139.3 ± 23.7	137.8 ± 23.0	0.607
DBP, mmHg	86.0 ± 16.2	84.7 ± 15.6	0.482
eGFR, mL/min/1.73 m ²	44.9 ± 36.5	43.3 ± 35.2	0.711
Albumin, g/dL	3.9 ± 0.5	3.9 ± 0.5	0.988
Hemoglobin, g/dL	12.3 ± 2.3	12.4 ± 1.9	0.637
UPCR, g/g Cr	0.8 ± 0.5	0.8 ± 0.6	0.654

Values are presented as mean ± SD or number (%).

HT-N, hypertensive nephrosclerosis; SBP, systolic blood pressure; DBP, diastolic blood pressure; eGFR, estimated glomerular filtration rate; UPCR, urine protein-creatinine ratio.