

Supplementary Table 4. Comparison of characteristics between subjects with eGFR < 60 mL/min/1.73 m² and eGFR ≥ 60 mL/min/1.73 m²

Variable	eGFR ≥ 60 mL/min/1.73 m ² (n = 1,368)	eGFR < 60 mL/min/1.73 m ² (n = 217)	p value
Age, yr	42.2 ± 11.7	52.6 ± 9.7	< 0.001
Male sex	618 (45.2)	127 (58.5)	< 0.001
Systolic blood pressure, mmHg	115.3 ± 27.1	115.1 ± 31.0	0.933
Body mass index, kg/m ²	23.8 ± 3.0	24.2 ± 2.7	0.056
< 18.5	25 (1.9)	5 (2.4)	
18.5–23	522 (39.9)	58 (28.3)	0.009
23–25	354 (27.1)	59 (28.8)	
≥ 25	407 (31.1)	83 (40.5)	
Hemoglobin, g/dL	13.7 ± 1.6	14.0 ± 1.5	0.033
Calcium, mg/dL	9.3 ± 0.4	9.3 ± 0.4	0.648
Phosphorus, mg/dL	3.5 ± 0.6	3.4 ± 0.5	0.018
Glucose, mg/dL	99.4 ± 16.8	102.5 ± 16.3	0.011
Uric acid, mg/dL	5.0 ± 1.4	5.3 ± 1.4	0.002
Cholesterol, mg/dL	187.4 ± 34.1	192.8 ± 31.6	0.027
Albumin, g/dL	4.4 ± 0.3	4.4 ± 0.3	0.808
eGFR, pre-donation, mL/min/1.73 m ²	101.3 ± 14.3	85.1 ± 12.9	< 0.001
eGFR, 1-month post-donation, mL/min/1.73 m ²	70.1 ± 14.1	53.1 ± 9.7	< 0.001
eGFR change from pre- to 1-month post-donation, %	30.8 ± 11.2	37.1 ± 10.3	< 0.001
Follow-up duration, mo	62.8 ± 49.4	58.2 ± 48.7	0.200

Values are presented as mean ± standard deviation or number (%).

eGFR, estimated glomerular filtration rate.