

Supplementary Table 2. MPN-SAF 10 symptoms within the last 12 months (patient reported)

Variable	MF		PV		ET		All	
	Score	Incidence	Score	Incidence	Score	Incidence	Score	Incidence
Fatigue or tiredness	4.89 ± 2.36	30 (76.92)	3.44 ± 2.41	13 (52.00)	5.00 ± 2.26	32 (78.05)	4.63 ± 2.39	75 (71.43)
Abdominal discomfort	2.33 ± 2.17	8 (20.51)	2.50 ± 3.02	2 (8.00)	3.07 ± 2.63	9 (21.95)	2.63 ± 2.48	19 (18.10)
Night sweats	3.38 ± 2.90	8 (20.51)	3.07 ± 2.89	9 (36.00)	1.80 ± 2.94	5 (12.20)	2.84 ± 2.90	22 (20.95)
Filling up quickly when eating (early satiety)	3.56 ± 2.71	10 (25.64)	2.67 ± 3.24	5 (20.00)	2.75 ± 3.02	4 (9.76)	3.08 ± 2.89	19 (18.10)
Bone pain (diffuse/not joint pain or arthritis)	1.67 ± 1.95	4 (10.26)	2.00 ± 1.95	7 (28.00)	1.67 ± 1.97	3 (7.32)	1.76 ± 1.91	14 (13.33)
Inactivity	3.54 ± 2.45	16 (41.03)	3.15 ± 2.61	8 (32.00)	3.77 ± 3.14	6 (14.63)	3.50 ± 2.64	30 (28.57)
Itching (pruritus)	2.75 ± 2.35	7 (17.95)	4.29 ± 3.24	15 (60.00)	2.70 ± 1.87	12 (29.27)	3.23 ± 2.58	34 (32.38)
Unintentional weight loss	3.50 ± 3.42	5 (12.82)	1.86 ± 3.76	0	1.58 ± 3.09	3 (7.32)	2.39 ± 3.38	8 (7.62)
Problems with concentration	3.83 ± 2.50	15 (38.46)	2.64 ± 2.84	8 (32.00)	2.81 ± 2.40	7 (17.07)	3.24 ± 2.55	30 (28.57)
Fever	2.08 ± 2.61	6 (15.38)	1.33 ± 2.18	3 (12.00)	1.30 ± 1.95	3 (7.32)	1.61 ± 2.25	12 (11.43)
MPN-SAF TSS								
Mean ± SD	16.84 ± 14.67		14.87 ± 10.15		13.59 ± 12.14		15.18 ± 12.73	
Median (range)	11.00 (0.00–56.00)		11.00 (2.00–36.00)		9.00 (0.00–53.00)		11.00 (0.00–56.00)	

Values are presented as mean ± standard deviation or number (%).

MPN-SAF, myeloproliferative neoplasm-Symptom Assessment Form; MF, myelofibrosis; PV, polycythemia vera; ET, essential thrombocythemia; TSS, Total Symptom Score; SD, standard deviation.