

Supplementary Table 2. The response rate for symptoms at 4 and 7 days after medication

	At 4 days after medication	At 7 days after medication
DW1601 (n = 67)	52 (78.8)	67 (100)
DW16011 (n = 70)	41 (60.3)	60 (85.7)
<i>P. sidoides</i> (n = 64)	34 (53.1)	55 (85.9)
Difference (95% CI) ^a	18.5 (3.25–33.74)	14.3 (6.09–22.48)
Difference (95% CI) ^b	25.7 (9.95–41.37)	14.1 (5.55–22.58)
<i>p</i> value ^c	0.020 ^a	0.001 ^a
<i>p</i> value ^d	0.002 ^a	0.002 ^a

Values are presented as number (%). The improvement rate for symptoms was defined as total bronchitis severity score (BSS) ≤ 3 or reduction of total BSS ≥ 7 .

P. sidoides, *Pelargonium sidoides*; CI, confidence interval.

^aThe difference was calculated as the value in the DW16011 group minus the value in the DW1601 group.

^bThe difference was calculated as the value in the *P. sidoides* group minus the value in the DW1601 group.

^c*p* value was calculated using Pearson's chi-square test for the difference of score between DW1601 and DW16011 groups.

^d*p* value was calculated using Pearson's chi-square test for the difference of score between DW1601 and *P. sidoides* groups.