

Supplementary Table 4. Incidence of metachronous gastric cancer according to characteristics at the time of endoscopic resection of gastric cancer

Variable	Total	Hanaoka et al. [21], 2010	Han et al. [19], 2011	Maehara et al. [26], 2012	Kim et al. [25], 2016	Moribata et al. [23], 2016	Oura et al. [27], 2018	Kato et al. [24], 2021
Location of gastric cancer								
Upper	30/202 (14.9)	-	3/21 (14.3)	2/25 (8.0)	6/34 (17.6)	2/25 (8.0)	1/23 (4.3)	-
Lower to mid	199/2,293 (8.7)	-	6/155 (3.9)	26/243 (10.7)	41/296 (13.9)	13/232 (5.6)	49/542 (9.0)	-
Gross type								64/422 (15.2)
Elevated	62/616 (10.1)	-	1/83 (1.2)	11/105 (10.5)	17/129 (13.2)	-	-	-
Depressed	95/569 (16.7)	-	6/21 (28.6)	17/163 (10.4)	30/201 (14.9)	-	-	42/184 (22.8)
Lauren's classification of cancer								
Intestinal type	75/809 (9.3)	-	-	27/259 (10.4)	-	-	48/550 (8.7)	-
Diffuse type	216 (12.5)	-	-	1/7 (14.3)	-	-	1/9 (11.1)	-
Differentiation of primary cancer								
Differentiated	62/798 (7.8)	-	-	-	-	14/249 (5.6)	48/549 (8.7)	-
Undifferentiated	3/24 (12.5)	-	-	-	-	1/8 (12.5)	2/16 (12.5)	-
Depth of invasion								
Mucosa	152/1,468 (10.4)	-	-	25/245 (10.2)	-	14/240 (5.8)	46/513 (9.0)	-
Submucosa	20/105 (19.0)	-	-	3/23 (13.0)	-	1/17 (5.9)	4/52 (7.7)	-
Synchronous gastric cancer								
Absent	77/436 (17.7)	-	-	-	-	-	-	77/436 (17.7)
Present	10/47 (21.3)	-	-	-	-	-	-	10/47 (21.3)
Curability of endoscopic treatment (eCura)								
Low risk	38/413 (9.2)	-	-	-	-	-	38/413 (9.2)	-
Intermediate risk	2/5 (40.0)	-	-	-	-	-	2/5 (40.0)	-
High risk	0 (0)	-	-	-	-	-	0 (0)	-

Values are presented as n/N (%).