

Supplementary Table 2. Baseline characteristics of RA patients with microbiologically confirmed NTM-PD

Variable	Value (n = 35)
Demographics	
Age, yr	66.2 ± 9.7
Sex	
Female	27 (77.1)
Male	8 (22.9)
BMI, kg/m ² (n = 32)	20.3 ± 3.1
Social history	
Smoking	6 (17.1)
Alcohol	3 (8.6)
Comorbidities	
Diabetes mellitus	7 (20.0)
Hypertension	13 (37.1)
Cardiovascular disease	5 (14.3)
Cancer	3 (8.6)
History of tuberculosis	7 (20.0)
ILD	8 (22.9)
COPD	5 (14.3)
Disease status at RA diagnosis	
RF positivity	28 (80.0)
RF titer, IU/mL	136.7 ± 193.8
Anti-CCP Ab positivity (n = 23)	18 (78.3)
Anti-CCP Ab titer, IU/mL (n = 23)	102.7 ± 110.2
Tender joint count	5.6 ± 6.8
Swollen joint count	5.8 ± 7.3
Patient global assessment, mm	46.4 ± 20.1
ESR, mm/h	44.4 ± 27.5
CRP, mg/dL	1.9 ± 2.9
DAS28	4.8 ± 1.7
Disease activity	
Low	4 (11.4)
Moderate	13 (37.1)
High	15 (42.9)
WBC, /μL	8,536.3 ± 3,322.8
Neutrophil, /μL (n = 34)	5,833.7 ± 2,814.1
Lymphocyte, /μL (n = 34)	1,957.8 ± 698.0
Hb, mg/dL	12.2 ± 1.5
Hct, %	36.8 ± 4.0
Platelet, × 10 ³ /μL	297.9 ± 108.1
Total protein, g/dL (n = 34)	7.3 ± 0.6
Albumin, g/dL (n = 34)	4.1 ± 0.4

Values are presented as mean ± standard deviation or number (%).

Anti-CCP Ab, anti-cyclic citrullinated peptide antibody; BMI, body mass index; COPD, chronic obstructive pulmonary disease; CRP, C-reactive protein; DAS, disease activity score; ESR, erythrocyte sedimentation rate; Hb, hemoglobin; Hct, hematocrit; ILD, interstitial lung disease; NTM-PD, non-tuberculous mycobacterial pulmonary disease; RA, rheumatoid arthritis; RF, rheumatoid factor; WBC, white blood cell.