

Supplementary Table 3. Initial *H. pylori* eradication rate based on endoscopic findings

Variable	Success (n = 198)	Failure (n = 30)	p value
M/F	126/72 (63.6)	17/13 (56.7)	0.752
Age (yr)	60.6 ± 10.3	62.1 ± 11.5	0.444
<i>H. pylori</i> IgG titer (U/mL)	4.6 ± 2.2	4.7 ± 2.3	0.877
Endoscopic morphologic features			
<i>H. pylori</i> non-infection			
RAC	2 (1.0)	0 (0.0)	> 0.999
Fundic gland polyp	2 (1.0)	0 (0.0)	> 0.999
Linear streak	1 (0.5)	0 (0.0)	> 0.999
Raised erosion	5 (2.5)	1 (3.3)	> 0.999
Hematin	5 (2.5)	1 (3.3)	> 0.999
<i>H. pylori</i> infection			
Atrophy type	194 (98.0)	30 (100.0)	0.969
Closed	48 (24.2)	12 (40.0)	0.154
Open	146 (73.7)	18 (60.0)	
Intestinal metaplasia	122 (61.6)	22 (75.9)	0.237
Diffuse redness	87 (43.9)	20 (66.7)	0.033
Spotty redness	152 (76.8)	22 (73.3)	0.856
Edema	30 (15.2)	7 (23.3)	0.386
Sticky mucus	53 (26.8)	14 (46.7)	0.044
Enlarged folds	14 (7.1)	2 (6.7)	> 0.999
Nodularity	16 (8.1)	1 (3.3)	0.583
Xanthoma	47 (23.7)	7 (23.3)	> 0.999
Hyperplastic polyp	19 (9.6)	1 (3.3)	0.433
<i>H. pylori</i> past infection			
Patchy redness	5 (2.5)	1 (3.3)	> 0.999
Map-like redness	5 (2.5)	0 (0.0)	0.833
Multiple flat white lesions	0 (0.0)	0 (0.0)	> 0.999

Values are presented as number (%) or mean ± standard deviation.

H. pylori, *Helicobacter pylori*; M, male; F, female; RAC, regular arrangement of collecting venules.