

Supplementary Table 2. Characteristics of the participants according to the presence or absence of typical chest pain

Characteristic	Patients without typical chest pain (n = 803)	Patients with typical chest pain (n = 3,652)	p value
SDT, day	1.983 ± 5.145	1.160 ± 7.588	< 0.001
Sex			< 0.001
Male	503 (62.6)	2,601 (71.2)	
Female	300 (37.4)	1,051 (23.6)	
Age, yr	68.96 ± 10.76	64.52 ± 11.63	< 0.001
Age ≥ 75 yr	275 (34.2)	802 (22.0)	< 0.001
EMS utilization	137 (17.1)	534 (14.6)	0.08
Final diagnosis			< 0.001
STEMI	235 (29.3)	1,716 (47.0)	
NSTEMI	568 (70.7)	1,936 (53.0)	
Killip functional class			< 0.001
I–II	502 (62.5)	3,159 (86.5)	
III–IV	301 (37.5)	493 (13.5)	
In-hospital death	86 (10.7)	132 (3.6)	< 0.001

Values are presented as number (percentage) for categorical values and mean ± standard deviation for continuous variables.

EMS, emergency medical service; NSTEMI, non-ST-elevation myocardial infarction; STEMI, ST-elevation myocardial infarction; SDT, symptom-to-door time.