

Supplementary Table 3. Clinical characteristics of patients with UIP and NSIP including linear regression model

	UIP (n = 32)	NSIP (n = 8)	<i>p</i> value
Age at sampling (yr)			
Mean $\pm$ SD	66.6 $\pm$ 8.7	65.3 $\pm$ 10.8	0.710
Median (IQR)	66.5 (13.5)	65.0 (17.5)	0.759
Age at diagnosis (yr)			
Mean $\pm$ SD	58.3 $\pm$ 11.3	56.6 $\pm$ 6.4	0.689
Median (IQR)	57.0 (17.8)	56.0 (8.5)	0.784
Sex, women	19 (59.4)	5 (62.5)	1.000
Smoking, ever	13 (40.6)	2 (25.0)	0.686
RF positivity	29 (90.6)	6 (75.0)	0.257
RF titer (IU/mL)			
Mean $\pm$ SD	261.2 $\pm$ 265.3	248.1 $\pm$ 219.5	0.904
Median (IQR)	171.6 (345.4)	189.0 (455.3)	0.929
Anti-CCP antibody	30 (93.8)	6 (85.7)	0.457
Anti-CCP antibody titer (U/mL)			
Mean $\pm$ SD	1,649.4 $\pm$ 1,868.1	1,525.7 $\pm$ 1,664.6	0.874
Median (IQR)	979.0 (1,700.9)	901.0 (2,478.0)	0.888
DAS28-ESR			
Mean $\pm$ SD	3.4 $\pm$ 1.1	4.0 $\pm$ 1.0	0.292
Median (IQR)	3.4 (1.2)	4.0 (1.7)	0.283
DAS28-CRP			
Mean $\pm$ SD	2.7 $\pm$ 1.0	2.8 $\pm$ 0.8	0.814
Median (IQR)	2.3 (1.8)	2.7 (1.4)	0.671
TL			
Mean $\pm$ SD	7.4 $\pm$ 1.9	10.3 $\pm$ 3.1	0.026
Median (IQR)	7.5 (2.4)	10.4 (5.0)	0.025

Values are presented as mean $\pm$ SD, median (IQR), or number (%).

UIP, usual interstitial pneumonia; NSIP, nonspecific interstitial pneumonia; SD, standard deviation; IQR, interquartile range; RF, rheumatoid factor; CCP, cyclic citrullinated peptide; DAS, disease activity score; ESR, erythrocyte sedimentation rate; CRP, C-reactive protein, TL, telomere length.

Patients with UIP and NSIP subtype were matched 1:1 by age and sex.