

Supplementary Table 1. Detailed information on antibiotic treatment

Variable	Total (n = 14)	Microbiological cure (n = 7)	Refractory (n = 7)
Macrolide	14 (100.0)	7 (100.0)	7 (100.0)
Duration (mo)	30 (16–40)	27 (13–38)	31 (26–76)
Ethambutol	9 (64.3)	5 (71.4)	4 (57.1)
Duration (mo)	14 (13–15)	13 (13–15)	14 (10–15)
Rifampicin/rifabutin	8 (57.1)	3 (42.8)	5 (71.4)
Duration (mo)	16 (13–20)	13 (12–16)	17 (15–23)
Fluoroquinolone	2 (14.3)	-	2 (28.6)
Duration (mo)	2 (1–3)	-	2 (1–3)
IV Amikacin	7 (50.0)	2 (28.6)	5 (71.4)
Duration (mo)	39 (25–102)	112 (39–184)	26 (25–52)
Inhaled amikacin	7 (50.0)	2 (28.6)	5 (71.4)
Duration (mo)	40 (20–93)	110 (40–181)	20 (20–53)
Imipenem/meropenem	7 (50.0)	2 (28.6)	5 (71.4)
Duration (mo)	2 (2–9)	2 (2–2)	6 (2–9)
Tigecycline	7 (50.0)	2 (28.6)	5 (71.4)
Duration (mo)	8 (4–19)	3 (0–6)	9 (8–19)
Clofazimine	7 (50.0)	2 (28.6)	5 (71.4)
Duration (mo)	34 (18–51)	27 (18–37)	34 (32–51)
Linezolid	2 (14.3)	0 (0.0)	2 (28.6)
Duration (mo)	3 (2–4)	-	3 (2–4)

Values are presented as number (%) or median (interquartile range).