

Supplementary Table 3. Comparison of sepsis treatments between dobutamine and no-dobutamine groups

	Before PSM			After PSM		
	DOBU (+) (N = 108)	DOBU (-) (N = 1,719)	<i>p</i> value	DOBU (+) (N = 105)	DOBU (-) (N = 209)	<i>p</i> value
Antibiotic adequacy	97 (89.8)	1,503 (87.4)	0.467	95 (90.5)	193 (92.3)	0.571
Vasopressin	69 (63.9)	567 (33.0)	< 0.001	66 (62.9)	130 (62.2)	0.910
Steroid	42 (38.9)	386 (22.5)	< 0.001	41 (39.0)	86 (41.1)	0.720
MV	72 (66.7)	740 (43.0)	< 0.001	69 (65.7)	139 (66.5)	0.889
CRRT	39 (36.1)	304 (17.7)	< 0.001	36 (34.3)	71 (34.0)	0.956
3-h sepsis bundle						
Lactate measure	105 (97.2)	1,634 (95.1)	0.308	102 (97.1)	201 (96.2)	0.757
Blood culture	68 (63.0)	1,225 (71.3)	0.066	67 (63.8)	134 (64.1)	0.958
Antibiotics	64 (59.3)	1,161 (67.5)	0.076	62 (59.0)	107 (51.2)	0.188
Fluid therapy	80 (74.1)	1,398 (81.3)	0.063	79 (75.2)	154 (73.7)	0.767
Vasopressors	67 (62.0)	1,099 (63.9)	0.691	65 (61.9)	134 (64.1)	0.701

Values are presented as number (%).

PSM, propensity score matching; DOBU, dobutamine; MV, mechanical ventilation; CRRT, continuous renal replacement therapy.